



All Products

About Constipation

Special Offers

Buy Now

Proud Sponsor of the Colon Cancer Alliance

Home » All Products » Dulcolax® Laxative Tablets

Dulcolax® Laxative Tablets

For predictable, overnight relief



*Among Stimulant Laxatives



Dulcolax® Laxative Tablets are clinically proven to relieve occasional constipation and irregularity overnight, so you can feel like yourself in the morning.

- Relieves constipation and irregularity overnight
- Comfort-coated tablets for gentle, predictable constipation relief

[Learn More](#)

[Buy Now](#)

[Drug Facts, FAQs](#)

★★★★★ 4.8 out of 5 stars.

4.8 Rating out of 5 stars.

[170 Reviews](#)

166 out of 167 (99%) reviewers recommend this product

[Write a review](#) . This action will open a modal dialog.

You may also like



★★★★★
★★★★★ 4.7 out of 5 stars.
(29)

Dulcolax® Pink™ Tablets for Women

[Buy Now](#)

[Learn More](#)



★★★★★
★★★★★ 4.8 out of 5 stars.
(68)

Dulcolax® Stool Softener

[Buy Now](#)

[Learn More](#)



★★★★★
★★★★★ 4.8 out of 5 stars.
(28)

Dulcolax® Laxative Suppositories

[Buy Now](#)

[Learn More](#)



★★★★★
★★★★★ 4.8 out of 5 stars.
(16)

DulcoEase® Pink™ Stool Softener

[Buy Now](#)

[Learn More](#)



Dulcolax® Laxative Suppositories with DulcoGlide™ Applicators

[Buy Now](#)

[Learn More](#)

[Learn More](#)

[Buy Now](#)

[Drug Facts](#)

[FAQs](#)

[Video](#)

[Reviews](#)

Dulcolax® Laxative Tablets

Drug Facts

Active ingredient (in each tablet)	Purpose
Bisacodyl USP 5 mg.....	Stimulant laxative

Use

- for temporary relief of occasional constipation and irregularity
- this product generally produces bowel movement in 6 to 12 hours

Warnings

Do not use if you cannot swallow without chewing

Ask a doctor before use if you have

- stomach pain, nausea or vomiting
- a sudden change in bowel habits that lasts more than 2 weeks

When using this product

- do not chew or crush tablet(s)
- do not use within 1 hour after taking an antacid or milk
- it may cause stomach discomfort, faintness and cramps

Stop use and ask a doctor if

- you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions take with a glass of water

adults and children 12 years of age and over	1 to 3 tablets in a single daily dose
children 6 to under 12 years of age	1 tablet in a single daily dose
children under 6 years of age	ask a doctor

Other information

- store at 20-25°C (68-77°F)
- protect from excessive humidity
- blister packaged for your protection. Do not use if individual seals are broken

Inactive ingredients: acacia, acetylated monoglyceride, carnauba wax, cellulose acetate phthalate, corn starch, dibutyl phthalate, docusate sodium, gelatin, glycerin, iron oxides, kaolin, lactose, magnesium stearate, methylparaben, pharmaceutical glaze, polyethylene glycol, povidone, propylparaben, Red No. 30 lake, sodium benzoate, sorbitan monooleate, sucrose, talc, titanium dioxide, white wax, Yellow No. 10 lake

Product of Italy

Manufactured in Mexico



[Home](#)
[Buy Now](#)
[Special Offers](#)
[Frequently Asked Questions](#)
[Contact Us](#)
[Terms of Use](#)
[Privacy Statement](#)
[Sitemap](#)
[References](#)

Copyright © 2013, Boehringer Ingelheim Pharmaceuticals, Inc. All rights reserved.